

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTASXS1656

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *
* *

NEWS 1		Web Page for STN Seminar Schedule - N. America
NEWS 2	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS 3	AUG 06	FSTA enhanced with new thesaurus edition
NEWS 4	AUG 13	CA/CAPplus enhanced with additional kind codes for granted patents
NEWS 5	AUG 20	CA/CAPplus enhanced with CAS indexing in pre-1907 records
NEWS 6	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS 7	AUG 27	USPATOLD now available on STN
NEWS 8	AUG 28	CAS REGISTRY enhanced with additional experimental spectral property data
NEWS 9	SEP 07	STN AnaVist, Version 2.0, now available with Derwent World Patents Index
NEWS 10	SEP 13	FORIS renamed to SOFIS
NEWS 11	SEP 13	INPADOCDB enhanced with monthly SDI frequency
NEWS 12	SEP 17	CA/CAPplus enhanced with printed CA page images from 1967-1998
NEWS 13	SEP 17	CAPplus coverage extended to include traditional medicine patents
NEWS 14	SEP 24	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS 15	OCT 02	CA/CAPplus enhanced with pre-1907 records from Chemisches Zentralblatt
NEWS 16	OCT 19	BEILSTEIN updated with new compounds
NEWS 17	NOV 15	Derwent Indian patent publication number format enhanced
NEWS 18	NOV 19	WPIX enhanced with XML display format
NEWS 19	NOV 30	ICSD reloaded with enhancements
NEWS 20	DEC 04	LINPADOCDB now available on STN

NEWS 21 DEC 14 BEILSTEIN pricing structure to change
 NEWS 22 DEC 17 USPATOLD added to additional database clusters
 NEWS 23 DEC 17 IMSDRUGCONF removed from database clusters and STN
 NEWS 24 DEC 17 DGENE now includes more than 10 million sequences
 NEWS 25 DEC 17 TOXCENTER enhanced with 2008 MeSH vocabulary in
 MEDLINE segment
 NEWS 26 DEC 17 MEDLINE and LMEDLINE updated with 2008 MeSH
 vocabulary
 NEWS 27 DEC 17 CA/CAPLUS enhanced with new custom IPC display
 formats
 NEWS 28 DEC 17 STN Viewer enhanced with full-text patent content
 from USPATOLD
 NEWS 29 JAN 02 STN pricing information for 2008 now available
 NEWS 30 JAN 16 CAS patent coverage enhanced to include exemplified
 prophetic substances

NEWS EXPRESS 19 SEPTEMBER 2007: CURRENT WINDOWS VERSION IS V8.2,
 CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
 AND CURRENT DISCOVER FILE IS DATED 19 SEPTEMBER 2007.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS LOGIN Welcome Banner and News Items
 NEWS IPC8 For general information regarding STN implementation
 of IPC 8

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN Customer
 agreement. Please note that this agreement limits use to scientific
 research. Use for software development or design or implementation
 of commercial gateways or other similar uses is prohibited and may
 result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:57:55 ON 16 JAN 2008

=> File Medline EMBASE Biosis Caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 15:58:15 ON 16 JAN 2008

FILE 'EMBASE' ENTERED AT 15:58:15 ON 16 JAN 2008
 Copyright (c) 2008 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 15:58:15 ON 16 JAN 2008
 Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 15:58:15 ON 16 JAN 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (1,4 or 1, 4) (2A) ((acetylgalactosaminy l transferase) or
acetylgalactosaminy ltransferase)
L1 446 (1,4 OR 1, 4) (2A) ((ACETYL GALACTOSAMINY L TRANSFERASE)
OR ACETYL
GALACTOSAMINY LTRANSFERASE)

=> s campylobacter
L2 38104 CAMPYLOBACTER

=> s l1 (10A) l2
L3 9 L1 (10A) L2

=> s l1 p l2
MISSING OPERATOR L1 P L2
The search profile that was entered contains terms or
nested terms that are not separated by a logical operator.

=> s l1 (p) l2
L4 20 L1 (P) L2

=> s l1 (l) l2
L5 20 L1 (L) L2

=> s l3 or l4 or l5
L6 20 L3 OR L4 OR L5

=> duplicate
ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove
ENTER L# LIST OR (END):l6
DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, BIOSIS, CAPLUS'
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n
PROCESSING COMPLETED FOR L6
L7 10 DUPLICATE REMOVE L6 (10 DUPLICATES REMOVED)

=> d l7 1-10 bib ab

L7 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2007:289252 CAPLUS
DN 146:496524
TI Structural characterization of Campylobacter jejuni
lipooligosaccharide
outer cores associated with Guillain-Barre and Miller Fisher
syndromes
AU Godschalk, Peggy C. R.; Kuijf, Mark L.; Li, Jianjun; St.
Michael, Frank;

Ang, C. Wim; Jacobs, Bart C.; Karwaski, Marie-France; Brochu, Denis;

Moterassed, Ali; Endtz, Hubert P.; van Belkum, Alex; Gilbert, Michel

CS Department of Medical Microbiology and Infectious Diseases, University

Medical Center, Rotterdam, 3015 GD, Neth.

SO Infection and Immunity (2007), 75(3), 1245-1254

CODEN: INFIBR; ISSN: 0019-9567

PB American Society for Microbiology

DT Journal

LA English

AB Mol. mimicry between lipooligosaccharides (LOS) of *Campylobacter jejuni*

and gangliosides in peripheral nerves plays a crucial role in the pathogenesis of *C. jejuni*-related Guillain-Barre syndrome (GBS).

We have

analyzed the LOS outer core structures of 26 *C. jejuni* strains associated

with GBS and its variant, Miller Fisher syndrome (MFS), by capillary

electrophoresis coupled with electrospray ionization mass spectrometry.

Sixteen out of 22 (73%) GBS-associated and all 4 (100%) MFS-associated strains

expressed LOS with ganglioside mimics. GM1a was the most prevalent

ganglioside mimic in GBS-associated strains (10/22, 45%), and in eight of

these strains, GM1a was found in combination with GD1a mimics. All seven

strains isolated from patients with ophthalmoplegia (GBS or MFS) expressed

disialylated (GD3 or GD1c) mimics. Three out of 22 GBS-associated strains

(14%) did not express sialylated ganglioside mimics because their LOS

locus lacked the genes necessary for sialylation. Three other strains

(14%) did not express ganglioside mimics because of frameshift mutations

in either the *cstII* sialyltransferase gene or the *cgtB* galactosyltransferase gene. It is not possible to determine if these mutations

were already present during *C. jejuni* infection. This is the first report

in which mass spectrometry combined with DNA sequence data were used to

infer the LOS outer core structures of a large number of neuropathy-associated

C. jejuni strains. We conclude that mol. mimicry between gangliosides and

C. jejuni LOS is the presumable pathogenic mechanism in most cases of C.

jejuni-related GBS. However, our findings suggest that in some cases,

other mechanisms may play a role. Further examination of the disease etiol. in

these patients is mandatory.

RE.CNT 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2005:497944 CAPLUS

DN 144:101691

TI Genomic diversity in Campylobacter jejuni: identification of C. jejuni

81-176-specific genes

AU Poly, Frederic; Threadgill, Deborah; Stintzi, Alain

CS Department of Veterinary Pathobiology, College of Veterinary Medicine,

Oklahoma State University, Stillwater, OK, 74078, USA

SO Journal of Clinical Microbiology (2005), 43(5), 2330-2338

CODEN: JCMIDW; ISSN: 0095-1137

PB American Society for Microbiology

DT Journal

LA English

AB Since the publication of the complete genomic sequence of Campylobacter

jejuni NCTC 11168 in Feb. 2000, evidence has been compiling that suggests

C. jejuni strains exhibit high genomic diversity. In order to investigate

this diversity, the unique genomic DNA sequences from a nonsequenced

Campylobacter strain, C. jejuni 81-176, were identified by comparison with

C. jejuni NCTC 11168 by using a shotgun DNA microarray approach.

Up to 63

kb of new chromosomal DNA sequences unique to this pathogen were obtained.

Eighty-six open reading frames were identified by the presence of uninterrupted coding regions encoding a min. of 40 amino acids.

In addition,

this study shows that the whole-plasmid shotgun microarray approach is

effective and provides a comprehensive coverage of DNA regions that differ

between two closely related genomes. The two plasmids harbored by this

Campylobacter strain, pTet and pVir, were also sequenced, with coverages

of 2.5- and 2.9-fold, resp., representing 72 and 92% of their complete

nucleotide sequences. The unique chromosomal genes encode proteins

involved in capsule and lipooligosaccharide biosynthesis, restriction and

modification systems, and respiratory metabolism. Several of these unique

genes are likely associated with C. jejuni 81-176 fitness and virulence.

Interestingly, the comparison of C. jejuni 81-176 unique genes with those

of C. jejuni ATCC 43431 revealed a single gene which encodes a probable

TraG-like protein. The product of this gene might be associated with the

mechanism of C. jejuni invasion into epithelial cells. In conclusion,

this study extends the repertoire of C. jejuni genes and thus will permit

the construction of a composite and more comprehensive microarray of C.

jejuni.

RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 10 MEDLINE on STN DUPLICATE 1

AN 2005392412 MEDLINE

DN PubMed ID: 16005859

TI Chemoenzymatic synthesis of
2-azidoethyl-ganglio-oligosaccharides GD3,
GT3, GM2, GD2, GT2, GM1, and GD1a.

AU Blixt Ola; Vasiliu Daniela; Allin Kirk; Jacobsen Nathan; Warnock
Dawn;

Razi Nahid; Paulson James C; Bernatchez Stephane; Gilbert Michel;
Wakarchuk Warren

CS Carbohydrate Synthesis and Protein Expression Core D, Consortium
for

Functional Glycomics, The Scripps Research Institute, Department
of

Molecular Biology, CB-248A, 92037 La Jolla, USA..

olablixt@scripps.edu

SO Carbohydrate research, (2005 Sep 5) Vol. 340, No. 12, pp.
1963-72.

Journal code: 0043535. ISSN: 0008-6215.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, NON-U.S. GOV'T)

LA English

FS Priority Journals

EM 200510

ED Entered STN: 31 Jul 2005

Last Updated on STN: 19 Oct 2005

Entered Medline: 18 Oct 2005

AB We have synthesized several ganglio-oligosaccharide structures using

glycosyltransferases from *Campylobacter jejuni*. The enzymes, alpha-(2-->3/8)-sialyltransferase (Cst-II), beta-(1-->4)-N-acetylgalactosaminyltransferase (CgtA), and beta-(1-->3)-galactosyltransferase (CgtB), were produced in

large-scale

fermentation from *Escherichia coli* and further characterized based on

their acceptor specificities. 2-Azidoethyl-glycosides corresponding to the

oligosaccharides of GD3

(alpha-D-Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->3)-

beta-D-Galp-(1-->4)-beta-D-Glcp-), GT3

(alpha-D-Neup5Ac-(2-->8)-alpha-D-

Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->3)-beta-D-Galp-(1-->4)-beta-D-Glcp-),

GM2

(beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-), GD2

(beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-), GT2

(beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->8)-

alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-), and

GM1

(beta-D-Galp-(1-->3)-beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-

D-Galp-(1-->4)-beta-D-Glcp-) were synthesized in high yields (gram-scale).

In addition, a mammalian alpha-(2-->3)-sialyltransferase (ST3Gal I) was

used to sialylate GM1 and generate GD1a

(alpha-D-Neup5Ac-(2-->3)-beta-D-

Galp-(1-->3)-beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-

(1-->4)-beta-D-Glcp-) oligosaccharide. We also cloned and expressed a rat

UDP-N-acetylglucosamine-4'epimerase (GalNAcE) in *E. coli* AD202 cells for

cost saving in situ conversion of less expensive UDP-GlcNAc to UDP-GalNAc.

L7 ANSWER 4 OF 10 MEDLINE on STN

DUPLICATE 2

AN 2005087616 MEDLINE

DN PubMed ID: 15716397

TI Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy.

AU Goodfellow John A; Bowes Tyrone; Sheikh Kazim; Odaka Masaaki; Halstead

Susan K; Humphreys Peter D; Wagner Eric R; Yuki Nobuhiro; Furukawa Koichi;

Furukawa Keiko; Plomp Jaap J; Willison Hugh J

CS Division of Clinical Neurosciences, Institute of Neurological Sciences,

Southern General Hospital, Glasgow G51 4TF, United Kingdom.

NC NS42888 (NINDS)

SO The Journal of neuroscience : the official journal of the Society for

Neuroscience, (2005 Feb 16) Vol. 25, No. 7, pp. 1620-8.

Journal code: 8102140. E-ISSN: 1529-2401.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

(RESEARCH SUPPORT, N.I.H., EXTRAMURAL)

(RESEARCH SUPPORT, NON-U.S. GOV'T)

(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)

LA English

FS Priority Journals

EM 200510

ED Entered STN: 19 Feb 2005

Last Updated on STN: 14 Oct 2005

Entered Medline: 13 Oct 2005

AB Anti-GD1a ganglioside antibodies (Abs) are the serological hallmark of the

acute motor axonal form of the post-infectious paralysis, Guillain-Barre

syndrome. Development of a disease model in mice has been impeded by the

weak immunogenicity of gangliosides and the apparent resistance of

GD1a-containing neural membranes to anti-GD1a antibody-mediated injury.

Here we used mice with altered ganglioside biosynthesis to generate such a

model at motor nerve terminals. First, we bypassed immunological tolerance by immunizing GD1a-deficient, beta-1,4-N-acetylgalactosaminyl transferase knock-out mice with GD1a ganglioside-mimicking antigens from Campylobacter jejuni and generated high-titer anti-GD1a antisera and complement fixing monoclonal Abs (mAbs). Next, we exposed ex vivo nerve-muscle preparations

from GD1a-overexpressing, GD3 synthase knock-out mice to the anti-GD1a

mAbs in the presence of a source of complement and investigated

morphological and electrophysiological damage. Dense antibody and complement deposits were observed only over presynaptic motor axons, accompanied by severe ultrastructural damage and electrophysiological blockade of motor nerve terminal function. Perisynaptic Schwann cells and postsynaptic membranes were unaffected. In contrast, normal mice were not only unresponsive to immunization with GD1a but also resistant to neural injury during anti-GD1a Ab exposure, demonstrating the central role of membrane antigen density in modulating both immune tolerance to GD1a and axonal susceptibility to anti-GD1a Abmediated injury. Identical paralyzing effects were observed when testing mouse and human anti-GD1a-positive sera. These data indicate that anti-GD1a Abs arise via molecular mimicry and are likely to be clinically relevant in injuring peripheral nerve axonal membranes containing sufficiently high levels of GD1a.

L7 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2002:276514 CAPLUS
 DN 136:320378
 TI Campylobacter glycosyltransferase genes and enzymes for biosynthesis of gangliosides and ganglioside mimics
 IN Gilbert, Michel; Wakarchuk, Warren W.
 PA National Research Council of Canada, Can.
 SO U.S. Pat. Appl. Publ., 84 pp., Cont.-in-part of U.S. Ser. No. 495,406.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.
DATE	-----	----	-----	-----
PI	US 2002042369	A1	20020411	US 2001-816028
20010321				
	US 6699705	B2	20040302	
	US 6503744	B1	20030107	US 2000-495406
20000131				
	EP 1652927	A2	20060503	EP 2005-25316
20000201				

EP 1652927	A3	20060719	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
	IE, SI, LT, LV, FI, RO, MK, CY, AL		
AT 329036	T	20060615	AT 2000-901455
20000201			
PT 1147200	T	20061031	PT 2000-901455
20000201			
ES 2269098	T3	20070401	ES 2000-901455
20000201			
CA 2441570	A1	20020926	CA 2002-2441570
20020222			
WO 2002074942	A2	20020926	WO 2002-CA229
20020222			
WO 2002074942	A3	20030313	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,			
CH, CN,			
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,		
GE, GH,			
	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,		
LK, LR,			
	LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,		
OM, PH,			
	PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR,		
TT, TZ,			
	UA, UG, US, UZ, VN, YU, ZA, ZM, ZW		
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,			
AZ, BY,			
	KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI,		
FR, GB,			
	GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI,		
CM, GA,			
	GN, GQ, GW, ML, MR, NE, SN, TD, TG		
AU 2002237122	A1	20021003	AU 2002-237122
20020222			
AU 2002237122	B2	20070322	
EP 1385941	A2	20040204	EP 2002-703414
20020222			
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
	IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		
JP 2004524033	T	20040812	JP 2002-574334
20020222			
US 2003148459	A1	20030807	US 2002-303161
20021121			
US 7138258	B2	20061121	
US 2003157655	A1	20030821	US 2002-303118
20021121			
US 6905867	B2	20050614	
US 2003157656	A1	20030821	US 2002-303128
20021121			

US 6911337	B2	20050628	
US 2003157657	A1	20030821	US 2002-303134
20021121			
US 6825019	B2	20041130	
US 2003157658	A1	20030821	US 2002-303162
20021121			
US 6723545	B2	20040420	
MX 2003PA08565	A	20040521	MX 2003-PA8565
20030922			
US 2004180406	A1	20040916	US 2003-735419
20031211			
US 7026147	B2	20060411	
US 2006166317	A1	20060727	US 2003-734719
20031211			
US 7169914	B2	20070130	
US 2004203103	A1	20041014	US 2004-820536
20040407			
US 7211657	B2	20070501	
US 2004229313	A1	20041118	US 2004-821573
20040408			
US 7192756	B2	20070320	
US 2004229263	A1	20041118	US 2004-821604
20040408			
US 2004265875	A1	20041230	US 2004-830825
20040424			
US 2004203112	A1	20041014	US 2004-845408
20040512			
US 7169593	B2	20070130	
US 2004203113	A1	20041014	US 2004-845412
20040512			
US 7166717	B2	20070123	
US 2004219638	A1	20041104	US 2004-846219
20040514			
US 7202353	B2	20070410	
US 2004229272	A1	20041118	US 2004-847983
20040517			
US 7208304	B2	20070424	
US 2004259203	A1	20041223	US 2004-850125
20040519			
US 7220848	B2	20070522	
US 2004259140	A1	20041223	US 2004-850807
20040521			
US 7217549	B2	20070515	
US 2005048630	A1	20050303	US 2004-962334
20041008			
US 7189836	B2	20070313	
US 2005084891	A1	20050421	US 2004-962235
20041008			
US 7238509	B2	20070703	
US 2005227248	A1	20051013	US 2004-961882
20041008			

US 7078207	B2	20060718	
US 2007048854	A1	20070301	US 2006-548514
20061011			
AU 2007202898	A1	20070712	AU 2007-202898
20070622			

PRAI US 1999-118213P	P	19990201
US 2000-495406	A2	20000131
EP 2000-901455	A3	20000201
US 2001-816028	A	20010321
AU 2002-237122	A3	20020222
WO 2002-CA229	W	20020222
US 2002-303118	A3	20021121
US 2002-303128	A1	20021121
US 2002-303134	A3	20021121
US 2004-821604	A3	20040408

AB This invention provides *Campylobacter jejuni* glycosyltransferases, including a bifunctional sialyltransferase that has both an α 2,3- and an α 2,8-activity. A β 1,4-GaINAc transferase and a β 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). In addnl. embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases. The enzymes may be used in preparation of gangliosides, lysogangliosides, and mimics of gangliosides and lysogangliosides. Thus, *C. jejuni* gene cstI α 2,3-sialyltransferase, gene cstII bifunctional α 2,3/ α 2,8-sialyltransferase, gene cgtA β - 1, 4-N-acetylgalactosaminyltransferase, and gene cgtB β -1,3-galactosyltransferase enzymes were used to prepare the carbohydrate portion of gangliosides GM1a, GM2, GM3, GD1a, GD3, and GT1a.

L7 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2002:37514 CAPLUS
 DN 137:16281

TI The genetic bases for the variation in the lipo-oligosaccharide of the mucosal pathogen, *Campylobacter jejuni*. Biosynthesis of sialylated ganglioside mimics in the core oligosaccharide
 AU Gilbert, Michel; Karwaski, Marie-France; Bernatchez, Stephane; Young, N.
 Martin; Taboada, Eduardo; Michniewicz, Joseph; Cunningham, Anna-Maria;

Wakarchuk, Warren W.
 CS Institute for Biological Sciences, National Research Council of
 Canada,
 Ottawa, ON, K1A 0R6, Can.
 SO Journal of Biological Chemistry (2002), 277(1), 327-337
 CODEN: JBCHA3; ISSN: 0021-9258
 PB American Society for Biochemistry and Molecular Biology
 DT Journal
 LA English
 AB The lipo-oligosaccharide (LOS) biosynthesis loci from 11
 Campylobacter
 jejuni strains expressing a total of 8 different ganglioside
 mimics in
 their LOS outer cores were compared. Based on the organization
 of the
 genes, the 11 corresponding loci could be classified into 3
 classes, with
 one of them being clearly an intermediate evolutionary step
 between the
 other two. Comparative genomics and expression of specific
 glycosyltransferases combined with in vitro activity assays
 allowed
 identification of ≥ 5 distinct mechanisms that allow C. jejuni to
 vary the structure of the LOS outer core as follows: (1)
 different gene
 complements; (2) phase variation because of homopolymeric
 tracts; (3) gene
 inactivation by the deletion or insertion of a single base
 (without phase
 variation); (4) single mutation leading to the inactivation of a
 glycosyltransferase; and (5) single or multiple mutations
 leading to
 "allelic" glycosyltransferases with different acceptor
 specificities. The
 differences in the LOS outer core structures expressed by the 11
 C. jejuni
 strains examined can be explained by one or more of these 5
 mechanisms.
 RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT
 L7 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2001:763199 CAPLUS
 DN 135:328761
 TI Campylobacter jejuni α 1,4-N-
 acetylgalactosaminyltransferase gene
 IN Endo, Tetsuo; Kakita, Shingo; Koizumi, Satoshi; Ozaki, Akio
 PA Kyowa Hakko Kogyo Co., Ltd., Japan
 SO PCT Int. Appl., 44 pp.
 CODEN: PIXXD2
 DT Patent

LA Japanese

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
------------	------	------	-----------------

DATE

-----	----	-----	-----
-------	------	-------	-------

PI WO 2001077337	A1	20011018	WO 2001-JP3111
------------------	----	----------	----------------

20010411

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE,
GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR,
LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
UZ, VN,

YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,
CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,
TR, BF,

BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
AU 2001046895 A5 20011023 AU 2001-46895

20010411

EP 1275721	A1	20030115	EP 2001-919887
------------	----	----------	----------------

20010411

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
MC, PT,

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

US 2004072324 A1 20040415 US 2003-257292
20030127

PRAI JP 2000-109150 A 20000411

WO 2001-JP3111 W 20010411

AB α 1,4-N-acetylgalactosaminyltransferase

(α 1,4-GalNAc transferase) of

Campylobacter jejuni, its gene, and use in biosynthetic
production of

GalNAc-containing complex carbohydrates, are disclosed. Complex
carbohydrates

having oligosaccharide containing galactose or
N-acetylgalactosamine (GalNAc)

at the reducing end are produced. A GalNAc-containing complex
carbohydrate

can be economically produced in a large amount by bringing the
microorganism

with the expression of the above enzyme, UDP-GalNAc, a receptor
complex

carbohydrate in an aqueous medium. Oligosaccharide moiety can
be lactose,

N-acetylactosamine, lacto-N-neotetraose, lacto-N-tetraose, para-lacto-N-neohexaose, Lewis X, or Lewis a. Production of GalNAc α 1-4Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc and GalNAc α 1-4Gal β 1-4Glc in E. coli transformed with C. jejuni α 1,4-GalNAc transferase, is described.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2000:553711 CAPLUS
DN 133:161277
TI Campylobacter glycosyltransferases for biosynthesis of gangliosides and ganglioside mimics
IN Gilbert, Michel; Wakarchuk, Warren W.
PA National Research Council of Canada, Can.
SO PCT Int. Appl., 120 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.
WO 2000046379	A1	20000810	WO 2000-CA86
20000201			
W:			AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, ZA, TJ, TM
RW:			GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
US 6503744	B1	20030107	US 2000-495406
20000131			
CA 2360205	A1	20000810	CA 2000-2360205
20000201			
EP 1147200	A1	20011024	EP 2000-901455
20000201			
EP 1147200	B1	20060607	
R:			AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, LT, LV, FI, RO, CY

JP 2002535992 T 20021029 JP 2000-597438
 20000201
 AU 772569 B2 20040429 AU 2000-22743
 20000201
 EP 1652927 A2 20060503 EP 2005-25316
 20000201
 EP 1652927 A3 20060719
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
 MC, PT,

IE, SI, LT, LV, FI, RO, MK, CY, AL
 AT 329036 T 20060615 AT 2000-901455
 20000201
 PT 1147200 T 20061031 PT 2000-901455
 20000201
 ES 2269098 T3 20070401 ES 2000-901455
 20000201
 MX 2001PA07853 A 20030925 MX 2001-PA7853
 20010801
 AU 2004203474 A1 20040826 AU 2004-203474
 20040729
 AU 2004203474 B2 20070920
 AU 2007202898 A1 20070712 AU 2007-202898

20070622
 PRAI US 1999-118213P P 19990201
 US 2000-495406 A 20000131
 EP 2000-901455 A3 20000201
 WO 2000-CA86 W 20000201
 AU 2002-237122 A3 20020222

AB This invention provides prokaryotic glycosyltransferases,
 including a
 bifunctional sialyltransferase that has both an α 2,3- and an
 α 2,8- activity. A β 1,4-GalNAc transferase and a
 β 1,3-galactosyltransferase are also provided by the invention,
 as are

other glycosyltransferases and enzymes involved in synthesis of
 lipooligosaccharide (LOS). The glycosyltransferases can be
 obtained from,

for example, Campylobacter species, including C. jejuni. In
 addnl.

embodiments, the invention provides nucleic acids that encode the
 glycosyltransferases, as well as expression vectors and host
 cells for

expressing the glycosyltransferases.

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 9 OF 10 MEDLINE on STN DUPLICATE 3
 AN 2001053031 MEDLINE
 DN PubMed ID: 11083778
 TI Sialylation of lipooligosaccharide cores affects immunogenicity
 and serum

resistance of *Campylobacter jejuni*.

AU Guerry P; Ewing C P; Hickey T E; Prendergast M M; Moran A P
CS Enteric Diseases Department, Naval Medical Research Center,
Silver Spring,

Maryland 20910, USA.. guerryp@nmrc.navy.mil

NC 1 RO1 A143559

SO Infection and immunity, (2000 Dec) Vol. 68, No. 12, pp. 6656-62.

Journal code: 0246127. ISSN: 0019-9567.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, NON-U.S. GOV'T)
(RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)

LA English

FS Priority Journals

EM 200012

ED Entered STN: 22 Mar 2001

Last Updated on STN: 22 Mar 2001

Entered Medline: 13 Dec 2000

AB Three genes involved in biosynthesis of the lipooligosaccharide
(LOS) core

of *Campylobacter jejuni* MSC57360, the type strain of the HS:1
serotype, whose structure mimics GM(2) ganglioside, have been
cloned and

characterized. Mutation of genes encoding proteins with
homology to a

sialyl transferase (cstII) and a putative N-acetylmannosamine
synthetase

(neuCl), part of the biosynthetic pathway of N-acetylneuraminic
acid

(NeuNAc), have identical phenotypes. The LOS cores of these
mutants

display identical changes in electrophoretic mobility, loss of
reactivity

with cholera toxin (CT), and enhanced immunoreactivity with a
hyperimmune

polyclonal antiserum generated against whole cells of *C. jejuni*
MSC57360.

Loss of sialic acid in the core of the neuCl mutant was
confirmed by fast

atom bombardment mass spectrometry. Mutation of a gene encoding
a

putative beta-1,4-N-acetylgalactosaminyltransferase

(Cgt) resulted in LOS cores intermediate in electrophoretic
mobility between that of wild type and the mutants lacking

NeuNAc, loss of

reactivity with CT, and a reduced immunoreactivity with
hyperimmune

antiserum. Chemical analyses confirmed the loss of
N-acetylgalactosamine

(GalNAc) and the presence of NeuNAc in the cgt mutant. These data suggest that the Cgt enzyme is capable of transferring GalNAc to an acceptor with or without NeuNAc and that the Cst enzyme is capable of transferring NeuNAc to an acceptor with or without GalNAc. A mutant with a nonsialylated LOS core is more sensitive to the bactericidal effects of human sera than the wild type or the mutant lacking GalNAc.

L7 ANSWER 10 OF 10 MEDLINE on STN DUPLICATE 4
AN 2000127862 MEDLINE
DN PubMed ID: 10660542
TI Biosynthesis of ganglioside mimics in *Campylobacter jejuni* OH4384.
Identification of the glycosyltransferase genes, enzymatic synthesis of model compounds, and characterization of nanomole amounts by 600-mhz (1)h and (13)c NMR analysis.
AU Gilbert M; Brisson J R; Karwaski M F; Michniewicz J; Cunningham A M; Wu Y; Young N M; Wakarchuk W W
CS Institute for Biological Sciences, National Research Council of Canada, Ottawa, Ontario K1A 0R6, Canada.
SO The Journal of biological chemistry, (2000 Feb 11) Vol. 275, No. 6, pp. 3896-906.
Journal code: 2985121R. ISSN: 0021-9258.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
OS GENBANK-AF130466; GENBANK-AF130984; GENBANK-AF167345
EM 200003
ED Entered STN: 27 Mar 2000
Last Updated on STN: 27 Mar 2000
Entered Medline: 16 Mar 2000
AB We have applied two strategies for the cloning of four genes responsible for the biosynthesis of the GT1a ganglioside mimic in the lipooligosaccharide (LOS) of a bacterial pathogen, *Campylobacter jejuni* OH4384, which has been associated with Guillain-Barre syndrome. We first cloned a gene encoding an alpha-2, 3-sialyltransferase (cst-I) using an activity screening strategy. We then used nucleotide sequence information from the recently completed sequence from *C. jejuni*
NCTC 11168

to amplify a region involved in LOS biosynthesis from *C. jejuni* OH4384.

The LOS biosynthesis locus from *C. jejuni* OH4384 is 11.47 kilobase pairs

and encodes 13 partial or complete open reading frames, while the corresponding locus in *C. jejuni* NCTC 11168 spans 13.49 kilobase pairs and

contains 15 open reading frames, indicating a different organization

between these two strains. Potential glycosyltransferase genes were

cloned individually, expressed in *Escherichia coli*, and assayed using

synthetic fluorescent oligosaccharides as acceptors. We identified genes

encoding a beta-1, 4-N-acetylgalactosaminyl-transferase (cgtA), a beta-1, 3-galactosyltransferase (cgtB), and a bifunctional sialyltransferase (cst-II), which transfers sialic acid to

O-3 of galactose and to O-8 of a sialic acid that is linked alpha-2,3- to

a galactose. The linkage specificity of each identified glycosyltransferase was confirmed by NMR analysis at 600 MHz on nanomole

amounts of model compounds synthesized in vitro. Using a gradient inverse

broadband nano-NMR probe, sequence information could be obtained by

detection of (3)J(C,H) correlations across the glycosidic bond. The role

of cgtA and cst-II in the synthesis of the GT1a mimic in *C. jejuni* OH4384

were confirmed by comparing their sequence and activity with corresponding

homologues in two related *C. jejuni* strains that express shorter ganglioside mimics in their LOS.

=> s l1 and l2

L8 28 L1 AND L2

=> duplicate

ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove

ENTER L# LIST OR (END):l8

DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, BIOSIS, CAPLUS'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n

PROCESSING COMPLETED FOR L8

L9 17 DUPLICATE REMOVE L8 (11 DUPLICATES REMOVED)

=> d l9 1-17 bib ab

L9 ANSWER 1 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN
AN 2007:455632 BIOSIS
DN PREV200700454271
TI beta 1,4-N-acetylgalactosaminyl transferases from C. jejuni.
AU Anonymous; Gilbert, Michel [Inventor]; Wakarchuk, Warren W. [Inventor]
CS Hull, Canada
ASSIGNEE: National Research Council of Canada
PI US 07238509 20070703
SO Official Gazette of the United States Patent and Trademark Office Patents,
(JUL 3 2007)
CODEN: OGUPE7. ISSN: 0098-1133.
DT Patent
LA English
ED Entered STN: 22 Aug 2007
Last Updated on STN: 22 Aug 2007
AB This invention provides prokaryotic glycosyltransferases, including a bifunctional sialyltransferase that has both an alpha 2,3- and an alpha 2,8-activity. A beta 1,4-GalNAc transferase and a beta 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). The glycosyltransferases can be obtained from, for example, Campylobacter species, including C. jejuni. In additional embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases.

L9 ANSWER 2 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN
AN 2007:231051 BIOSIS
DN PREV200700230171
TI Nucleic acids encoding beta 1,4-N-acetylgalactosaminyltransferases from C. jejuni.
AU Anonymous; Gilbert, Michel [Inventor]; Wakarchuk, Warren W. [Inventor]
CS Hull, Canada
ASSIGNEE: National Research Council of Canada
PI US 07189836 20070313
SO Official Gazette of the United States Patent and Trademark Office Patents,
(MAR 13 2007)
CODEN: OGUPE7. ISSN: 0098-1133.

DT Patent
LA English
ED Entered STN: 4 Apr 2007
Last Updated on STN: 4 Apr 2007
AB This invention provides prokaryotic glycosyltransferases, including a bifunctional sialyltransferase that has both an alpha 2,3- and an alpha 2,8-activity. A beta 1,4-GalNAc transferase and a beta 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). The glycosyltransferases can be obtained from, for example, Campylobacter species, including C. jejuni. In additional embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases.

L9 ANSWER 3 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN
AN 2007:185941 BIOSIS
DN PREV200700192247
TI beta 1,4-N-acetylgalactosaminyltransferases from C. jejuni.
AU Anonymous; Gilbert, Michel [Inventor]; Wakarchuk, Warren W. [Inventor]
CS Hull, Canada
ASSIGNEE: National Research Council of Canada
PI US 07169593 20070130
SO Official Gazette of the United States Patent and Trademark Office Patents,
(JAN 30 2007)
CODEN: OGUPE7. ISSN: 0098-1133.

DT Patent
LA English
ED Entered STN: 14 Mar 2007
Last Updated on STN: 14 Mar 2007
AB This invention provides prokaryotic glycosyltransferases, including a bifunctional sialyltransferase that has both an alpha 2,3- and an alpha 2,8-activity. A beta 1,4-GalNAc transferase and a beta 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). The glycosyltransferases can be obtained from, for example, Campylobacter species, including C. jejuni. In additional embodiments, the invention provides nucleic acids that encode

the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases.

L9 ANSWER 4 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN
AN 2007:117784 BIOSIS
DN PREV200700116026
TI Nucleic acids encoding beta 1,4-N-acetylgalactosaminyltransferases from C. jejuni.
AU Anonymous; Gilbert, Michel [Inventor]; Wakarchuk, Warren W. [Inventor]
CS Hull, Canada
ASSIGNEE: National Research Council of Canada
PI US 07166717 20070123
SO Official Gazette of the United States Patent and Trademark Office Patents,
(JAN 23 2007)
CODEN: OGUPE7. ISSN: 0098-1133.
DT Patent
LA English
ED Entered STN: 14 Feb 2007
Last Updated on STN: 14 Feb 2007
AB This invention provides prokaryotic glycosyltransferases, including a bifunctional sialyltransferase that has both an alpha 2,3- and an alpha 2,8-activity. A beta 1,4-GalNAc transferase and a beta 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). The glycosyltransferases can be obtained from, for example, Campylobacter species, including C. jejuni. In additional embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases.

L9 ANSWER 5 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2007:289252 CAPLUS
DN 146:496524
TI Structural characterization of Campylobacter jejuni lipooligosaccharide outer cores associated with Guillain-Barre and Miller Fisher syndromes
AU Godschalk, Peggy C. R.; Kuijff, Mark L.; Li, Jianjun; St. Michael, Frank;
Ang, C. Wim; Jacobs, Bart C.; Karwaski, Marie-France; Brochu, Denis;

Moterassed, Ali; Endtz, Hubert P.; van Belkum, Alex; Gilbert, Michel

CS Department of Medical Microbiology and Infectious Diseases, University

Medical Center, Rotterdam, 3015 GD, Neth.

SO Infection and Immunity (2007), 75(3), 1245-1254

CODEN: INFIBR; ISSN: 0019-9567

PB American Society for Microbiology

DT Journal

LA English

AB Mol. mimicry between lipooligosaccharides (LOS) of *Campylobacter jejuni* and gangliosides in peripheral nerves plays a crucial role in the

pathogenesis of *C. jejuni*-related Guillain-Barre syndrome (GBS).

We have

analyzed the LOS outer core structures of 26 *C. jejuni* strains associated

with GBS and its variant, Miller Fisher syndrome (MFS), by capillary

electrophoresis coupled with electrospray ionization mass spectrometry.

Sixteen out of 22 (73%) GBS-associated and all 4 (100%) MFS-associated strains

expressed LOS with ganglioside mimics. GM1a was the most prevalent

ganglioside mimic in GBS-associated strains (10/22, 45%), and in eight of

these strains, GM1a was found in combination with GD1a mimics.

All seven

strains isolated from patients with ophthalmoplegia (GBS or MFS) expressed

disialylated (GD3 or GD1c) mimics. Three out of 22 GBS-associated strains

(14%) did not express sialylated ganglioside mimics because their LOS

locus lacked the genes necessary for sialylation. Three other strains

(14%) did not express ganglioside mimics because of frameshift mutations

in either the *cstII* sialyltransferase gene or the *cgtB* galactosyltransferase gene. It is not possible to determine if these mutations

were already present during *C. jejuni* infection. This is the first report

in which mass spectrometry combined with DNA sequence data were used to

infer the LOS outer core structures of a large number of neuropathy-associated

C. jejuni strains. We conclude that mol. mimicry between gangliosides and

C. jejuni LOS is the presumable pathogenic mechanism in most cases of *C.*

jejuni-related GBS. However, our findings suggest that in some cases,

other mechanisms may play a role. Further examination of the disease etiol. in

these patients is mandatory.

RE.CNT 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 6 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson
Corporation on STN

AN 2006:235396 BIOSIS

DN PREV200600241377

TI Nucleic acids encoding beta-1,4-GalNAc transferase.

AU Gilbert, Michel [Inventor]; Wakarchuk, Warren W. [Inventor]

CS ASSIGNEE: National Research Council of Canada

PI US 06911337 20050628

SO Official Gazette of the United States Patent and Trademark
Office Patents,

(JUN 28 2005)

CODEN: OGUPE7. ISSN: 0098-1133.

DT Patent

LA English

ED Entered STN: 19 Apr 2006

Last Updated on STN: 19 Apr 2006

AB This invention provides prokaryotic glycosyltransferases,
including a

bifunctional sialyltransferase that has both an alpha 2,3- and
an alpha

2,8-activity. A beta 1,4-GalNAc transferase and a beta

1,3-galactosyltransferase are also provided by the invention, as
are other

glycosyltransferases and enzymes involved in synthesis of
lipooligosaccharide (LOS). The glycosyltransferases can be
obtained from,

for example, Campylobacter species, including C. jejuni. In

additional embodiments, the invention provides nucleic acids
that encode

the glycosyltransferases, as well as expression vectors and host
cells for

expressing the glycosyltransferases.

L9 ANSWER 7 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2005:497944 CAPLUS

DN 144:101691

TI Genomic diversity in Campylobacter jejuni: identification of C.
jejuni 81-176-specific genes

AU Poly, Frederic; Threadgill, Deborah; Stintzi, Alain

CS Department of Veterinary Pathobiology, College of Veterinary
Medicine,

Oklahoma State University, Stillwater, OK, 74078, USA

SO Journal of Clinical Microbiology (2005), 43(5), 2330-2338

CODEN: JCMIDW; ISSN: 0095-1137

PB American Society for Microbiology

DT Journal

LA English

AB Since the publication of the complete genomic sequence of *Campylobacter jejuni* NCTC 11168 in Feb. 2000, evidence has been compiling that suggests *C. jejuni* strains exhibit high genomic diversity.

In order to investigate this diversity, the unique genomic DNA sequences

from a nonsequenced *Campylobacter* strain, *C. jejuni* 81-176, were identified by comparison with *C. jejuni* NCTC 11168 by using a shotgun DNA

microarray approach. Up to 63 kb of new chromosomal DNA sequences unique

to this pathogen were obtained. Eighty-six open reading frames were

identified by the presence of uninterrupted coding regions encoding a min.

of 40 amino acids. In addition, this study shows that the whole-plasmid

shotgun microarray approach is effective and provides a comprehensive

coverage of DNA regions that differ between two closely related genomes.

The two plasmids harbored by this *Campylobacter* strain, pTet and pVir, were also sequenced, with coverages of 2.5- and 2.9-fold, resp.,

representing 72 and 92% of their complete nucleotide sequences. The

unique chromosomal genes encode proteins involved in capsule and lipooligosaccharide biosynthesis, restriction and modification systems,

and respiratory metabolism. Several of these unique genes are likely associated

with *C. jejuni* 81-176 fitness and virulence. Interestingly, the comparison of *C. jejuni* 81-176 unique genes with those of *C. jejuni* ATCC

43431 revealed a single gene which encodes a probable TraG-like protein.

The product of this gene might be associated with the mechanism of *C. jejuni*

invasion into epithelial cells. In conclusion, this study extends the

repertoire of *C. jejuni* genes and thus will permit the construction of a

composite and more comprehensive microarray of *C. jejuni*.

RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

AN 2005392412 MEDLINE
 DN PubMed ID: 16005859
 TI Chemoenzymatic synthesis of
 2-azidoethyl-ganglio-oligosaccharides GD3,
 GT3, GM2, GD2, GT2, GM1, and GD1a.
 AU Blixt Ola; Vasiliu Daniela; Allin Kirk; Jacobsen Nathan; Warnock
 Dawn;
 Razi Nahid; Paulson James C; Bernatchez Stephane; Gilbert Michel;
 Wakarchuk Warren
 CS Carbohydrate Synthesis and Protein Expression Core D, Consortium
 for
 Functional Glycomics, The Scripps Research Institute, Department
 of
 Molecular Biology, CB-248A, 92037 La Jolla, USA..
 olablixt@scripps.edu
 SO Carbohydrate research, (2005 Sep 5) Vol. 340, No. 12, pp.
 1963-72.
 Journal code: 0043535. ISSN: 0008-6215.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 LA English
 FS Priority Journals
 EM 200510
 ED Entered STN: 31 Jul 2005
 Last Updated on STN: 19 Oct 2005
 Entered Medline: 18 Oct 2005
 AB We have synthesized several ganglio-oligosaccharide structures
 using
 glycosyltransferases from *Campylobacter jejuni*. The enzymes,
 alpha-(2-->3/8)-sialyltransferase (Cst-II), beta-(1-->4
)-N-acetylgalactosaminyltransferase (CgtA), and
 beta-(1-->3)-galactosyltransferase (CgtB), were produced in
 large-scale
 fermentation from *Escherichia coli* and further characterized
 based on
 their acceptor specificities. 2-Azidoethyl-glycosides
 corresponding to the
 oligosaccharides of GD3
 (alpha-D-Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->3)-
 beta-D-Galp-(1-->4)-beta-D-Glcp-), GT3
 (alpha-D-Neup5Ac-(2-->8)-alpha-D-
 Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->3)-beta-D-Galp-(1-->4)-beta-D-Glc
 p-),
 GM2
 (beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-
 beta-D-Glcp-), GD2
 (beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->8)-alpha-
 D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-), GT2
 (beta-D-GalpNAc-(1-->4)-[alpha-D-Neup5Ac-(2-->8)-alpha-D-Neup5Ac-(2-->
 8)-

alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-), and GM1 (beta-D-Galp-(1-->3)-beta-D-GalpNac-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-) were synthesized in high yields (gram-scale). In addition, a mammalian alpha-(2-->3)-sialyltransferase (ST3Gal I) was used to sialylate GM1 and generate GD1a (alpha-D-Neup5Ac-(2-->3)-beta-D-Galp-(1-->3)-beta-D-GalpNac-(1-->4)-[alpha-D-Neup5Ac-(2-->3)]-beta-D-Galp-(1-->4)-beta-D-Glcp-) oligosaccharide. We also cloned and expressed a rat UDP-N-acetylglucosamine-4'epimerase (GalNacE) in E. coli AD202 cells for cost saving in situ conversion of less expensive UDP-GlcNac to UDP-GalNac.

L9 ANSWER 9 OF 17 MEDLINE on STN DUPLICATE 2
 AN 2005087616 MEDLINE
 DN PubMed ID: 15716397
 TI Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy.
 AU Goodfellow John A; Bowes Tyrone; Sheikh Kazim; Odaka Masaaki; Halstead Susan K; Humphreys Peter D; Wagner Eric R; Yuki Nobuhiro; Furukawa Koichi; Furukawa Keiko; Plomp Jaap J; Willison Hugh J
 CS Division of Clinical Neurosciences, Institute of Neurological Sciences, Southern General Hospital, Glasgow G51 4TF, United Kingdom.
 NC NS42888 (NINDS)
 SO The Journal of neuroscience : the official journal of the Society for Neuroscience, (2005 Feb 16) Vol. 25, No. 7, pp. 1620-8. Journal code: 8102140. E-ISSN: 1529-2401.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE) (RESEARCH SUPPORT, N.I.H., EXTRAMURAL) (RESEARCH SUPPORT, NON-U.S. GOV'T) (RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)
 LA English
 FS Priority Journals
 EM 200510
 ED Entered STN: 19 Feb 2005

Last Updated on STN: 14 Oct 2005

Entered Medline: 13 Oct 2005

AB Anti-GD1a ganglioside antibodies (Abs) are the serological hallmark of the acute motor axonal form of the post-infectious paralysis, Guillain-Barre syndrome. Development of a disease model in mice has been impeded by the weak immunogenicity of gangliosides and the apparent resistance of GD1a-containing neural membranes to anti-GD1a antibody-mediated injury.

Here we used mice with altered ganglioside biosynthesis to generate such a model at motor nerve terminals. First, we bypassed immunological tolerance by immunizing GD1a-deficient, beta-1,4-N-acetylgalactosaminyl transferase knock-out mice with GD1a ganglioside-mimicking antigens from Campylobacter jejuni and generated high-titer anti-GD1a antisera and complement fixing monoclonal Abs (mAbs). Next, we exposed ex vivo nerve-muscle preparations from GD1a-overexpressing, GD3 synthase knock-out mice to the anti-GD1a mAbs in the presence of a source of complement and investigated morphological and electrophysiological damage. Dense antibody and complement deposits were observed only over presynaptic motor axons, accompanied by severe ultrastructural damage and electrophysiological blockade of motor nerve terminal function. Perisynaptic Schwann cells and postsynaptic membranes were unaffected. In contrast, normal mice were not only unresponsive to immunization with GD1a but also resistant to neural injury during anti-GD1a Ab exposure, demonstrating the central role of membrane antigen density in modulating both immune tolerance to GD1a and axonal susceptibility to anti-GD1a Abmediated injury. Identical paralyzing effects were observed when testing mouse and human anti-GD1a-positive sera. These data indicate that anti-GD1a Abs arise via molecular mimicry and are likely to be clinically relevant in injuring peripheral nerve axonal membranes containing sufficiently high levels of GD1a.

AN 2002:276514 CAPLUS
 DN 136:320378
 TI Campylobacter glycosyltransferase genes and enzymes for
 biosynthesis of gangliosides and ganglioside mimics
 IN Gilbert, Michel; Wakarchuk, Warren W.
 PA National Research Council of Canada, Can.
 SO U.S. Pat. Appl. Publ., 84 pp., Cont.-in-part of U.S. Ser. No.
 495,406.

CODEN: USXXCO

DT Patent
 LA English
 FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.
US 2002042369	A1	20020411	US 2001-816028
US 6699705	B2	20040302	
US 6503744	B1	20030107	US 2000-495406
EP 1652927	A2	20060503	EP 2005-25316
EP 1652927	A3	20060719	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
AT 329036	T	20060615	AT 2000-901455
PT 1147200	T	20061031	PT 2000-901455
ES 2269098	T3	20070401	ES 2000-901455
CA 2441570	A1	20020926	CA 2002-2441570
WO 2002074942	A2	20020926	WO 2002-CA229
WO 2002074942	A3	20030313	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,			

	KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI,		
FR, GB,			
	GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI,		
CM, GA,			
	GN, GQ, GW, ML, MR, NE, SN, TD, TG		
AU 2002237122	A1	20021003	AU 2002-237122
20020222			
AU 2002237122	B2	20070322	
EP 1385941	A2	20040204	EP 2002-703414
20020222			
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,		
MC, PT,			
	IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		
JP 2004524033	T	20040812	JP 2002-574334
20020222			
US 2003148459	A1	20030807	US 2002-303161
20021121			
US 7138258	B2	20061121	
US 2003157655	A1	20030821	US 2002-303118
20021121			
US 6905867	B2	20050614	
US 2003157656	A1	20030821	US 2002-303128
20021121			
US 6911337	B2	20050628	
US 2003157657	A1	20030821	US 2002-303134
20021121			
US 6825019	B2	20041130	
US 2003157658	A1	20030821	US 2002-303162
20021121			
US 6723545	B2	20040420	
MX 2003PA08565	A	20040521	MX 2003-PA8565
20030922			
US 2004180406	A1	20040916	US 2003-735419
20031211			
US 7026147	B2	20060411	
US 2006166317	A1	20060727	US 2003-734719
20031211			
US 7169914	B2	20070130	
US 2004203103	A1	20041014	US 2004-820536
20040407			
US 7211657	B2	20070501	
US 2004229313	A1	20041118	US 2004-821573
20040408			
US 7192756	B2	20070320	
US 2004229263	A1	20041118	US 2004-821604
20040408			
US 2004265875	A1	20041230	US 2004-830825
20040424			
US 2004203112	A1	20041014	US 2004-845408
20040512			
US 7169593	B2	20070130	

US 2004203113	A1	20041014	US 2004-845412
20040512			
US 7166717	B2	20070123	
US 2004219638	A1	20041104	US 2004-846219
20040514			
US 7202353	B2	20070410	
US 2004229272	A1	20041118	US 2004-847983
20040517			
US 7208304	B2	20070424	
US 2004259203	A1	20041223	US 2004-850125
20040519			
US 7220848	B2	20070522	
US 2004259140	A1	20041223	US 2004-850807
20040521			
US 7217549	B2	20070515	
US 2005048630	A1	20050303	US 2004-962334
20041008			
US 7189836	B2	20070313	
US 2005084891	A1	20050421	US 2004-962235
20041008			
US 7238509	B2	20070703	
US 2005227248	A1	20051013	US 2004-961882
20041008			
US 7078207	B2	20060718	
US 2007048854	A1	20070301	US 2006-548514
20061011			
AU 2007202898	A1	20070712	AU 2007-202898
20070622			
PRAI US 1999-118213P	P	19990201	
US 2000-495406	A2	20000131	
EP 2000-901455	A3	20000201	
US 2001-816028	A	20010321	
AU 2002-237122	A3	20020222	
WO 2002-CA229	W	20020222	
US 2002-303118	A3	20021121	
US 2002-303128	A1	20021121	
US 2002-303134	A3	20021121	
US 2004-821604	A3	20040408	

AB This invention provides *Campylobacter jejuni* glycosyltransferases, including a bifunctional sialyltransferase that has both an α 2,3- and an α 2,8-activity. A β 1,4-GaINAc transferase and a β 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). In addnl. embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the

glycosyltransferases. The enzymes may be used in preparation of gangliosides, lysogangliosides, and mimics of gangliosides and lysogangliosides. Thus,

C. jejuni gene cstI α 2,3-sialyltransferase, gene cstII bifunctional

α 2,3/ α 2,8-sialyltransferase, gene cgtA β - 1,

4-N-acetylgalactosaminyltransferase, and gene cgtB

β -1,3-galactosyltransferase enzymes were used to prepare the carbohydrate portion of gangliosides GM1a, GM2, GM3, GD1a, GD3, and GT1a.

L9 ANSWER 11 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN

AN 2003:30396 BIOSIS

DN PREV200300030396

TI Synthesis of oligosaccharides and glycolipids using bacterial glycosyltransferases.

AU Johnson, Karl F. [Reprint Author]; Bezila, Dan [Reprint Author]; Gbewonyo,

Hugh [Reprint Author]; Ngo, Winnie [Reprint Author]; Garber, Colby

[Reprint Author]; Taylor, Diane E.

CS Neose Technologies, Inc., Horsham, PA, USA

SO Glycobiology, (October 2002) Vol. 12, No. 10, pp. 706. print. Meeting Info.: 7th Annual Conference of the Society for Glycobiology.

Boston, MA, USA. November 09-12, 2002. Society for Glycobiology. ISSN: 0959-6658.

DT Conference; (Meeting)

Conference; (Meeting Poster)

Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 8 Jan 2003

Last Updated on STN: 8 Jan 2003

L9 ANSWER 12 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN

AN 2003:30393 BIOSIS

DN PREV200300030393

TI Evolution of lipopolysaccharide glycosyltransferase specificity examined

by the study of homologous enzymes.

AU Wakarchuk, Warren W. [Reprint Author]; Bernatchez, Stephane [Reprint

Author]; Gilbert, Michel [Reprint Author]; Karwaski, Marie-France [Reprint

Author]; Masson, Amara [Reprint Author]; Logan, Susan [Reprint Author];

Dunn, J  ssica [Reprint Author]
CS Institute for Biological Sciences, National Research Council of
Canada,

100 Sussex Drive, Ottawa, ON, K1A 0R6, Canada

SO Glycobiology, (October 2002) Vol. 12, No. 10, pp. 705-706.
print.

Meeting Info.: 7th Annual Conference of the Society for
Glycobiology.

Boston, MA, USA. November 09-12, 2002. Society for Glycobiology.

ISSN: 0959-6658.

DT Conference; (Meeting)

Conference; (Meeting Poster)

Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 8 Jan 2003

Last Updated on STN: 11 Feb 2003

L9 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2002:37514 CAPLUS

DN 137:16281

TI The genetic bases for the variation in the lipo-oligosaccharide
of the

mucosal pathogen, *Campylobacter jejuni*. Biosynthesis of
sialylated ganglioside mimics in the core oligosaccharide

AU Gilbert, Michel; Karwaski, Marie-France; Bernatchez, Stephane;
Young, N.

Martin; Taboada, Eduardo; Michniewicz, Joseph; Cunningham,
Anna-Maria;

Wakarchuk, Warren W.

CS Institute for Biological Sciences, National Research Council of
Canada,

Ottawa, ON, K1A 0R6, Can.

SO Journal of Biological Chemistry (2002), 277(1), 327-337

CODEN: JBCHA3; ISSN: 0021-9258

PB American Society for Biochemistry and Molecular Biology

DT Journal

LA English

AB The lipo-oligosaccharide (LOS) biosynthesis loci from 11
Campylobacter jejuni strains expressing a total of 8 different
ganglioside mimics in their LOS outer cores were compared.

Based on the

organization of the genes, the 11 corresponding loci could be
classified

into 3 classes, with one of them being clearly an intermediate
evolutionary step between the other two. Comparative genomics
and

expression of specific glycosyltransferases combined with in
vitro

activity assays allowed identification of ≥ 5 distinct mechanisms
that allow *C. jejuni* to vary the structure of the LOS outer core

as

follows: (1) different gene complements; (2) phase variation because of homopolymeric tracts; (3) gene inactivation by the deletion or insertion of a single base (without phase variation); (4) single mutation leading to the inactivation of a glycosyltransferase; and (5) single or multiple mutations leading to "allelic" glycosyltransferases with different acceptor specificities. The differences in the LOS outer core structures expressed by the 11 C. jejuni strains examined can be explained by one or more of these 5 mechanisms.

RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2001:763199 CAPLUS
DN 135:328761
TI Campylobacter jejuni α 1,4-N-acetylgalactosaminyltransferase gene
IN Endo, Tetsuo; Kakita, Shingo; Koizumi, Satoshi; Ozaki, Akio
PA Kyowa Hakko Kogyo Co., Ltd., Japan
SO PCT Int. Appl., 44 pp.
CODEN: PIXXD2
DT Patent
LA Japanese
FAN.CNT 1

DATE	PATENT NO.	KIND	DATE	APPLICATION NO.
-----	-----	----	-----	-----

PI	WO 2001077337	A1	20011018	WO 2001-JP3111
20010411				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,			

BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 AU 2001046895 A5 20011023 AU 2001-46895
 20010411
 EP 1275721 A1 20030115 EP 2001-919887
 20010411

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
 MC, PT,

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
 US 2004072324 A1 20040415 US 2003-257292
 20030127

PRAI JP 2000-109150 A 20000411
 WO 2001-JP3111 W 20010411

AB α 1,4-N-acetylgalactosaminyltransferase
 (α 1,4-GalNAc transferase) of

Campylobacter jejuni, its gene, and use in biosynthetic
 production of

GalNAc-containing complex carbohydrates, are disclosed. Complex
 carbohydrates

having oligosaccharide containing galactose or
 N-acetylgalactosamine (GalNAc)

at the reducing end are produced. A GalNAc-containing complex
 carbohydrate

can be economically produced in a large amount by bringing the
 microorganism

with the expression of the above enzyme, UDP-GalNAc, a receptor
 complex

carbohydrate in an aqueous medium. Oligosaccharide moiety can
 be lactose,

N-acetyllactosamine, lacto-N-neotetraose, lacto-N-tetraose,
 para-lacto-N-neohexaose, Lewis X, or Lewis a. Production of
 GalNAc α 1-4Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc and
 GalNAc α 1-4Gal β 1-4Glc in E. coli transformed with C. jejuni
 α 1,4-GalNAc transferase, is described.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 15 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2000:553711 CAPLUS

DN 133:161277

TI Campylobacter glycosyltransferases for biosynthesis of
 gangliosides and ganglioside mimics

IN Gilbert, Michel; Wakarchuk, Warren W.

PA National Research Council of Canada, Can.

SO PCT Int. Appl., 120 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE			

PI WO 2000046379	A1	20000810	WO 2000-CA86
20000201			
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,			
CR, CU,			
CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,			
ID, IL,			
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,			
LV, MA,			
MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE,			
SG, ZA,			
TJ, TM			
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH,			
CY, DE,			
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF,			
BJ, CF,			
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6503744	B1	20030107	US 2000-495406
20000131			
CA 2360205	A1	20000810	CA 2000-2360205
20000201			
EP 1147200	A1	20011024	EP 2000-901455
20000201			
EP 1147200	B1	20060607	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
IE, LT, LV, FI, RO, CY			
JP 2002535992	T	20021029	JP 2000-597438
20000201			
AU 772569	B2	20040429	AU 2000-22743
20000201			
EP 1652927	A2	20060503	EP 2005-25316
20000201			
EP 1652927	A3	20060719	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
IE, SI, LT, LV, FI, RO, MK, CY, AL			
AT 329036	T	20060615	AT 2000-901455
20000201			
PT 1147200	T	20061031	PT 2000-901455
20000201			
ES 2269098	T3	20070401	ES 2000-901455
20000201			
MX 2001PA07853	A	20030925	MX 2001-PA7853
20010801			
AU 2004203474	A1	20040826	AU 2004-203474
20040729			
AU 2004203474	B2	20070920	
AU 2007202898	A1	20070712	AU 2007-202898
20070622			
PRAI US 1999-118213P	P	19990201	
US 2000-495406	A	20000131	

EP 2000-901455 A3 20000201
WO 2000-CA86 W 20000201
AU 2002-237122 A3 20020222

AB This invention provides prokaryotic glycosyltransferases, including a bifunctional sialyltransferase that has both an α 2,3- and an α 2,8- activity. A β 1,4-GalNAc transferase and a β 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). The glycosyltransferases can be obtained from, for example, Campylobacter species, including C. jejuni. In addnl. embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases.

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 16 OF 17 MEDLINE on STN DUPLICATE 3
AN 2001053031 MEDLINE
DN PubMed ID: 11083778
TI Sialylation of lipooligosaccharide cores affects immunogenicity and serum resistance of Campylobacter jejuni.
AU Guerry P; Ewing C P; Hickey T E; Prendergast M M; Moran A P
CS Enteric Diseases Department, Naval Medical Research Center, Silver Spring, Maryland 20910, USA.. guerryp@nmrc.navy.mil
NC 1 RO1 A143559
SO Infection and immunity, (2000 Dec) Vol. 68, No. 12, pp. 6656-62.

Journal code: 0246127. ISSN: 0019-9567.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 (RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)
 (RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)

LA English
FS Priority Journals
EM 200012

ED Entered STN: 22 Mar 2001
 Last Updated on STN: 22 Mar 2001
 Entered Medline: 13 Dec 2000

AB Three genes involved in biosynthesis of the lipooligosaccharide (LOS) core of Campylobacter jejuni MSC57360, the type strain of the HS:1 serotype, whose structure mimics GM(2) ganglioside, have been cloned and

characterized. Mutation of genes encoding proteins with homology to a sialyl transferase (cstII) and a putative N-acetylmannosamine synthetase (neuCl), part of the biosynthetic pathway of N-acetylneuraminic acid (NeuNAc), have identical phenotypes. The LOS cores of these mutants display identical changes in electrophoretic mobility, loss of reactivity with cholera toxin (CT), and enhanced immunoreactivity with a hyperimmune polyclonal antiserum generated against whole cells of *C. jejuni* MSC57360. Loss of sialic acid in the core of the neuCl mutant was confirmed by fast atom bombardment mass spectrometry. Mutation of a gene encoding a putative beta-1,4-N-acetylgalactosaminyltransferase (Cgt) resulted in LOS cores intermediate in electrophoretic mobility between that of wild type and the mutants lacking NeuNAc, loss of reactivity with CT, and a reduced immunoreactivity with hyperimmune antiserum. Chemical analyses confirmed the loss of N-acetylgalactosamine (GalNAc) and the presence of NeuNAc in the cgt mutant. These data suggest that the Cgt enzyme is capable of transferring GalNAc to an acceptor with or without NeuNAc and that the Cst enzyme is capable of transferring NeuNAc to an acceptor with or without GalNAc. A mutant with a nonsialylated LOS core is more sensitive to the bactericidal effects of human sera than the wild type or the mutant lacking GalNAc.

L9 ANSWER 17 OF 17 MEDLINE on STN DUPLICATE 4
 AN 2000127862 MEDLINE
 DN PubMed ID: 10660542
 TI Biosynthesis of ganglioside mimics in *Campylobacter jejuni* OH4384. Identification of the glycosyltransferase genes, enzymatic synthesis of model compounds, and characterization of nanomole amounts by 600-mhz (1)h and (13)c NMR analysis.
 AU Gilbert M; Brisson J R; Karwaski M F; Michniewicz J; Cunningham A M; Wu Y;
 Young N M; Wakarchuk W W
 CS Institute for Biological Sciences, National Research Council of Canada,

Ottawa, Ontario K1A 0R6, Canada.

SO The Journal of biological chemistry, (2000 Feb 11) Vol. 275, No. 6, pp.

3896-906.

Journal code: 2985121R. ISSN: 0021-9258.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

OS GENBANK-AF130466; GENBANK-AF130984; GENBANK-AF167345

EM 200003

ED Entered STN: 27 Mar 2000

Last Updated on STN: 27 Mar 2000

Entered Medline: 16 Mar 2000

AB We have applied two strategies for the cloning of four genes responsible

for the biosynthesis of the GT1a ganglioside mimic in the lipooligosaccharide (LOS) of a bacterial pathogen, *Campylobacter jejuni* OH4384, which has been associated with Guillain-Barre syndrome. We

first cloned a gene encoding an alpha-2, 3-sialyltransferase (cst-I) using

an activity screening strategy. We then used nucleotide sequence information from the recently completed sequence from *C. jejuni* NCTC 11168

to amplify a region involved in LOS biosynthesis from *C. jejuni* OH4384.

The LOS biosynthesis locus from *C. jejuni* OH4384 is 11.47 kilobase pairs

and encodes 13 partial or complete open reading frames, while the corresponding locus in *C. jejuni* NCTC 11168 spans 13.49 kilobase pairs and

contains 15 open reading frames, indicating a different organization

between these two strains. Potential glycosyltransferase genes were

cloned individually, expressed in *Escherichia coli*, and assayed using

synthetic fluorescent oligosaccharides as acceptors. We identified genes

encoding a beta-1, 4-N-acetylgalactosaminyl-transferase (cgtA), a beta-1, 3-galactosyltransferase (cgtB), and a bifunctional sialyltransferase (cst-II), which transfers sialic acid to

O-3 of galactose and to O-8 of a sialic acid that is linked alpha-2,3- to

a galactose. The linkage specificity of each identified glycosyltransferase was confirmed by NMR analysis at 600 MHz on nanomole

amounts of model compounds synthesized in vitro. Using a gradient inverse

broadband nano-NMR probe, sequence information could be obtained by

detection of $(3)J(C,H)$ correlations across the glycosidic bond. The role

of cgtA and cst-II in the synthesis of the GT1a mimic in *C. jejuni* OH4384

were confirmed by comparing their sequence and activity with corresponding

homologues in two related *C. jejuni* strains that express shorter ganglioside mimics in their LOS.

SEQ ID NO: 1

Aaagaatacgaatttgctaaagagg (41; would anneal to the non-coding strand)

ttttaaatcttagtggttatgtagaaccacatatagaattagcgccaaaatttaattcttgaagagctaattggcttttacaaaaatgatggatcttat
cataggaatgatagcgttccaacacatttagcttttgcctttaataaagcatctattacgatttttgggtcaacaccaagctaccgcaatgctttt
caaaactcatatcaataaaatcattgatacaggtaaaaaaatccaaatgccagcatatcgataaaagtgtattttgtatcacgcgtatagaagaag
aagatatcttcaaaacttgccaaaggcttacttaatgaaaaatagtgatagaatatatcttagtctttattatattttgaaattttttgttactttta
tgcctgattgtatcttgcattttttagctttgattgtagcaagaatcgcttttcatcttaacaaaaaacccgcgaataatcatcaatacaaaatttgca
aatctgttttccctcaatacactcttcaaaaagaacgcgataaaattgtcttttaaaaatttatgaaaaatttgctcaatttgggtattgattgttgc
caaaacaccaccaagaaaaaattctcaataaagttaaatttcatcaatgaaaattttcttatagatgcctggctttaaagcgtcctattatcttca
caactgcacactatggaaactgggaaattttaagccttgcttatgctggctaaatattggtgcgatttccatagtgaggaaaaaaggttaaaaagtgaagt
tatgtatgaaattttaagccaaagtgcacccaatttgacatagaacttattgacaaaaaggcggtataagacaaatgctaagtgtcttaaaaaag
gagagagctttgggaagcttttaactgtatgcaagactgcgtagaaaaaagaaagcgttaagattaaaaatttttaacaaagaagtgaattatcaaatgggag
caagccttatcgacacaaagaagcaatgctttgatcatcctgtttatgcctataaagaaggtggtaaattttgcatagagttttttaagcaaaaga
ttctcaaaatgcaagtttgaagaactgacactttatcaagcacaaagtgcgaagaaatgattaaaaaaagaccttgggaatacttttttttctat
agacgctttgtctagtataatgaggaatttacaaggggtcaaaaatgaatctaaaaacaaataagcgttattatcatcgtaaaaaatgctgagcaaac
tttgcctttgagttttaaattttaaagatttttgcaagatttgcgtagaaaaaatttatcttaaaaaatttatgaaaaatttgattgaagcattattcttaaggaatcaaatgggag
aaaaaagattttgtcaatttatatatatttatcaccaatgcttttataggttttggagcttttaaaaaatcttgctttaagttatgcaaaaaatgatggga
ttttaagcattgatgctgatgaagtgccttgaagaatgagtgatttaagagccttaaaaaatttaaaacttcaagaagataatcatcgcacttagccg
taaaaaatctctataaaggcgaatggataaaggcattgtggttgggtggcctgattatgttttgagaatttttaataaaaaatttcaactcgcttttaaatgat
aattttagtaacatgaaagccttttgccttgcgaagtaattgcgtagaaaaaatttatcttaaaaaatttgattgaagcattattcttaaggaatcaaatgggag
taattgacaaaaatgcagtaactactcaagctcttgggcaaaaacaaatatacacaaaaaaagtggtgttttaaaagcaaatttaagagctttttggac
tttttttagaaatttttttaaaaaatggccttttatatggtttataagggttttataattagcgtttgtctgcattgggaacattttttaaatat
atgaaatttatgaacttcaaaagcaaaaaacaaaaacttgcgcttttaataataaacttataatcaaaaaagaacgccttaaaactagtgcttgata
gtgttaaaacttgccttttaccacaaatgaagttttaaactgcagatggttgtaaaacttgcgttaagcatttattgaagcatttattgaagcatttcaaaaga
tttctcctgtccttttaaaacacatttggcaagaagatgaaggggtttaaacttagtaaaagtcgcaacaaaactataaaaaacgctgatagtgaaat
ataatagttattgatggtgatgatgtttggaaaaagatttcataaaagaacatttagaatttgacacaaagaagcgtttttttacaaggttcaagag
taatttttaataaaaaagaagcgaagaattttaacaaagatgattatcgcataaatttttaataaaaaagatttttaaaagtctcaaaaaattctt
tttagctaaaaatttttcaactcttcaaaaaaaagatgaaaaaattcttttaaaaaacactcttattaaaggtattaggggttgcaaatgatgttt
ttttaaactgatttttgtaacttgatggttttaattgaaattttattggttggggtagagaagatagtgaaatttgggtgctgattttttattta
aaaggcatttttagacgattaaaaatttaaagctattgcttatcatatttatcacaagaagaaatagcaaaaaatgcttgaaagcaatcatcaaat
atttagataccatcaaaaaataaaagatttctggagatataaacatgaagaaatagggttagttataccaatctataatgtgaaaaaatatttaag
agaatggttagatagcgttatcaatcaacttataactataacttagaaatcatacttgccttaagtgatggttagcacagatgaacactcaactcaatattgca
aaagaatatcacttcaaaagataaaagaataaactcttttgataaagaaaaatgggggtttaaagtcagatgaataataggatagaactcttagcg
gggaatataaattaaaaaacaaaactcaacatataaaagaaaattctttaaatagaatttcaattggatggttaataatccttataatataataaagc
atataaaagctctcaagcttttaataatgaaaaagatttaaccaattttacttacctagtagatttatatttatctttagatagtgataatatt
tggaactaaactgcataagaagaatgcgttataagaatgaaaaatgtggatgtattgtggttgaccatgattgcacattgaagacaataaaaa
attaaagcaaaaaaaacaggatggaatttttgatttttaaaaaagaatgtataatcactccaaaaagaatatgcaaatcgagcattaaagttagggtc
tagagatatcttttggatggaatggaatgattgtatttttaatttttaaaagcaaattaaacttaattttataaattttattatcaatgaagatata
cactttgggataattttgtttgctagtgctaaataaatttatgttttatcacaagaagttgtatttgtgtcgtttaagagcaaacagtagtatcaaat
atgataagaagattacaaaagcaaatgtgtcagagatttttaagatatatatgaaacttccggggaacgcctaaggaagcaaaaaattattttaa
agcaatatagcagggttaaaagcttttaaaatttgatagaatttttaaaagattcacaacaaacgaaaaatgcacttgctataaaaaagaacatttttacc
tgctatgccaaaaaagcctttaatgattaaaaaatttaaaaaagatccttttaatttaaaaggaacaaattagttttaattaaaccttttattcaaa
aacttctctatgatatttggaaattttggcaaaaaaataaaaaatatttaataataaaaaatataaaaaatataatttttaggtataatcactata
attataggagaaaatttttatatgctatttcaatcatacttttggtaaaaaatatttgccttattcatccttttagaaaaatttagacataaaaa
aaaacatttttaataaaaaacatacaacagagataaaatcgattcttataccaaaaaaactccttggtgcaaatataaatcaacataaaggaat
taattaaacttaataaagctattatagggggaggggcataaaggatattttaattatgatgaaaaatctaaagatccaaatctccttgaatccttg
ggcttttatacgagtaaaaaatgaagctattaccttaaaagcttctcttgaaagcatattgcctgctatccaaagaggtgttataggtataatgat
tgtaccgatggaagtgaagaataattctagaattttgcaaacatattcttcatttatccaataaaaaatctcttatgaatttcaaatcaaaac
caaaatcagaagaaaaataaactctatagctattataaatttatgttttgcaagttttatcccaaaagatgagtggttcttaaatagatggatcatat
ctatgatgctaaaaaactttataaaagcttctatataccaaaaaacaaatgatgtagtttagttattcaaggggttgatattcactatttttaatgat
aatttttttcttggtaagataaatatggcaatatattgaagaaccaggagattgcttgccttatcaataattataacttaaaatggaaagaagtat
taattgacagaatcaatacaaatggaaaaaagcaacaaaaaagttttcttcaaatatacactctttagagcaattaaagtataaacacaggaat
attatttccactgaattaaataatttatcattttcttttttaaaaaaacatagactcaagatatattataaatataattggatagaactattgaagaa
tttaaaaaattctattttacaaaatattatcatataaaatagaaccttctatgatttcaaaagaaactctaaaaaaaatattcttaacattgttttaa
aatttttttatatttaataaaaaatttttaaggttaaaatatttttttagctaaataatgttaaccattattttgttcttttttttatatatattgaa
tatatagcaaatattttaatttagcacatagagaacgctacaatacttgttttaaaaaataattttgccttaaatagtttttaaaaaacaaactgcaactcttg
aatattatttttaacaaagccttcaattcttagtattacaaatgttaatttattaggcacgtaaatgatataaattacagttcatatgctattttt
tgagcttgacttaacattggataatatacaatacatcttcagccatattgattttaacatcttctcgcagctttaaactcgcaaaagcttctaaat
acaatttctttctataagtttccccacatagtcacataaaattttcttggcaataattttttttacaaactctttttgctataaaaaaccaga
attaaagtcaaaactttttatgaaataacattactttcaacaatagcattgaaaaacactaaatcaacttcatctcgttcaataaaattttata
cactcttcacaagcatttagttccaataatcatcaggatctaaaaacattataaaggagagtttgctactttcacacctcatatgcttgcctta
aaagacctaaagtttttctgattgttttcttctttttagatattttttagatattttttagctatattttatctatttatcatttccacaatc
atcaactacaattatttctatatcttttaaaagctgattgatacagcttctattgcccctgctatattgttccacattataagttggtaagatg
attgaaatttttaacatatatttctctatttttatataatttaattataacataaaatctattttgataaaatcggttaaaaaatcaatcttgatgga
aaataacatgaaaaaagtattattgctggaaatggacaaagttaaaagaatttgatttcaagactaccaaatgattttgtatgtatttagatg
taactcaatttttttgaagataaatactatcttggtaaaaaatgcgaagcagatttttaacatctattcttttttgaacaatactacacttta
aaacatttaatacaaaatcaagaatagagaccgaactaattatgtgttctaattacaaccaagctcatctagaaaatgaaaaatttgtaaaaactt
tttacgattatttctgtagctcatttgggatagattttttcaaacacttaagatttttaatgcttatttttaatttcaagaaatttatttcaa
tcaagaattacctcaggggtctatattgtgtgcagtagccatagccctaggatcaaaagaatttatctttcgggaattgatttttatcaaatggg

tcattctttagcttttgataactaaacaaaaaatcttttaaaattggctcctaattttaaaaatgataattcacactatatcgacatagtaaaaaa
cagatataaaagcttttagaattttctagaaaaaacttacaaaaataaaactatattgcttattgtcctaacagctcttttagcaaaattttatagaactagc
gccaaattttaattcaaaattttatcatacaagaaaaaataaactacactaaagatatactcatatcttctagtgaggctttaggaaaaattttcaaaa
aatattaattttaaaaaataaaaaataaaagaaaatatttattacaagtgtataaaagatctattaagattaccttagtgatataaaagcattatttca
aaggaaaaataaatgaaagaaaataaaaaataaaataataatcataagtgaaagaaaaagcacccttagtcgctgctgaaataggcatttaataatgga
cagtttagaactagctaaaaattatggtagatgcagccttttagcacaggtgctaagattataaagcatcaaacccacatcgttgaagatgagatgagt
aaggccgctaaaaaagtaattcctgggtaattgcaaaaaataagcatttatgagattatgcaaaaaatgtgcttttagattataaaagatgagctagcactta
aagaatacacagaaaaattaggtcttgtttatcttagcacaccttttctcgtgcagggtgcaaacgcgttagaagatatgggaggttagtgcttttaa
gattgggttcaggtgagttgtaataattatccgcttattaaacacatagcagcctttaaaaaagccttagatagtttagcacagggatgaatagttgaa
agtataaaaaccaactgtaaaaatcttattagacaatgaaattccctttgttttaatgcacacaaccaatctttaccaaccccgcatatcttgttaa
gattaaacgctatgcttgaattaaaaaaagaattttcttgtatggtaggcttaagcgaccacacacagataatcttgcgtgttttaggtgcggttgc
acttgggtgcttgtgtgcttgaagacatttttactgatagtagcatagaagtgccctgatagtttgttctatggatcacacaggttttaaaagag
cttatttatacaaaatggctataatgagagcattataaaaaatgaaagttaaaaaagccttagatagtttagcacagggatgaatagttgcttttg
caagcgtagtcagcattaaagatatataaaaaaggcgaagttttatctatggataatatttgggttaaaagacctggacttgggtggaatttagtcagc
tgaatttgaataatttttaggcataaaagcatttaagagatatagaaaatgatactcagtttaagctatgaggatttttgcgtgaaaaaaatccttttta
taacaggcactagggtgatttattctaagattaaatctttaatgtacaggtgcaaaactcaagcgaatttgaactttacatcttttgaacaggaat
gcacttaagcaaaaattttggctatacagtttaagaactttataaaaaatggcttaaaaaatatttatgaattttataaattacgataaatttttca
accgataaggcttttagccatcacatattgatggattttcaagatatgttaattgagctaaaacctgatttaactcgtagtacatggagatagaatcgagc
ctttagcagcagctattgttggagcattaaacaatatcttagtagcacatattgaaggtggagagatttcaggaaactattgatgatagcttacgcca
cgctatatcaaaactagcacatattcatttagtaaatgatgagtttgcataaaaggcgtttaatgcagcttgggagaagatgaaaaatctatttttatc
ataggttcgccttagaacttttaaacgataataaaaaatctcacttataagtaagcaaaaaaataattatgatataaattatgaaaaactacgctttgc
ttattgtttctatcctgtttacaactgaaattacaagcattaaaaataaagcagatatttagtaaaagcactgatacaaaagtaacaaaaattatattgt
tatttatccaaataatgatttaggttttgaattaatcttgcataagctatgaagaacttaaaaaataaccctagatttaagcttttccatcgcttaga
tttgagtttttataaacttgttaaaaaatgctgattttataataggtaattcaagttgtattttaaaagaggccttatacttaaaaacagcaggaa
tttagttggctcaaggcaaaatggaagacttggcaatgaaaatacactaaaagtttaagcaaatagtgatgaaatactaaaagctatttaaacct
tcataaaaaacaagattttattttagcgccaagtttagagatttttagatagctcaaaaattatttttgaattttacaaagcggagaaatttttaaaaa
aacacacaaaaagtttttaaggatataaaatgagcttagcaataatccctgctcgtggtggctcaaaagggtattaaaaataaaaaatttgggtttatt
aaacaataaaccttttaatttattacaccattaaagctgcactaaatactaaaagcatttagtaaaagtgttgaagcagtgatagtgatgaaatttta
aattatgcaaaaagtcaaaatgttgatatttttaaacgcccatttagccttgcacagataatactacaagcgataaagtgcttttacatgctctaa
aattttacaaaagattatgagatagtttttttacaaccaccctcgccgctaagaacaaatattcatattgataggcttttaacttttataaaaa
tagcaatgcaaatgccctaatttagcgtaagcgaatgtgataataaaattctaaaagcctttgtttgtaatgaatatggcgatttagcagggtttgt
aatgatgaatatccttttatgccaaaggcaaaaattgcctaaaacatatatgagcaatggtgcaatttatatttttaagataaaagaatttttaaaaa
atcctagctttttacaaagcaaaaaccaagcattttttaaaggatgaaagctcaagtttagatattgactgtttggaggatttaaaaaaggctgaaca
gatattggaaaaaatcccttaaaatgcaataaaaaataatttaaatttattaagcaatataatatttatacaaaaacttatatagaaaaatcctagaa
gattttcaagactaaaaaccaagattttataacctttccatttgaaaaacaatcaactagagagtgtagcggggctggggatagaagaatattgtgc
ttttaaatttagcaatatcttatacagaaatggattcattttcttttagcgatcttttctacctcattatacaaaaagttggaaggtattgttcaatt
cttgatgggggtttctatgttttaactttcaacatcctatggatagaatcagcactgcaagttttacctatgaacaaaatcatagttttattaacgatg
cttgccaaaatcacatcaacaaaacatttctatagtttaacataatccaagctcatcaataacgcatttaattatacaagatgagtttggatagg
aaaagatgttttgcttaaacagggtatcacacttgggactggatgtgtcataggacaaagagctgtagttactaaagatgtaccaccttatgctata
gttgcaggaaattccagccaaaattatcaaatatagattttagtgaaaaaacaatagaagattattaaaaattcaatgggtggaatatcattttgctg
atttttatgatattgatctttaaatttaaaaaataaccaatatcttgacctactagaagaaaaaatcataaaaaaatcaatttctactataatccaaa
taaaactttatttttagagatttttagaactaaaatcaaaaaaatttttaactatttttaactctatttttcacccctgcttctctctcttttaaaac
ttcaataattttctgatgaaattcatcatgtgcaaaactctttggatagtttttttatgatttcattacttttctttttatcatgataattttgattt
aaaattttctttatttttattctcatatcttccatttggattaaattcataatgataaatgcaagtttttaaaaacagctattttctcacaaaacataa
ataaaatataacaaaaaagcacatcttcgccataattcaaacgctcatctattttaattttttcaaaaactttttaagatgatatttttttaaaagca
cttcgccccaaaccgaccgcaaaaatgcctttgttgccttaaaaaattctaaaaattccttttgattaaaaacttcacttctgttttaaaacgataaaaat
tgtttgtttttttaccctatgcacaaaggcatcaaaacaaagcaaatcaaaacctttttcatctcttttaaacgctatttcacaagcatcaggtgtta
aaaaatcatcactatctaaaaacattataaaatcagaactagaatgcaaaacccccaaatttctacttgcaaaagtgccataattttcttcattttg
aaagattttttattcttggatcttttttgcataattctaaacacatatataaactatttctttacttttatcatcgataatcaaaatttcaatatct
tttaagcttgattatacaacttgcataagctcttgagataaaatcgcaagaatttaaaaagcgggattatgatagaagttgtggcatatttttcc
taaaattttgttaaaataataaaaaaattctatcaagtttaggaaatttatgaaaatttttatcaccttccaa
cctggttaggcgatacggtaattggc (40, reverse complement; would anneal to the coding strand)

SEQ ID NO: 40.

gccattaccgtatgcctaaccagg

rev complement: cctggttaggcgatacggtaattggc

SEQ ID NO: 41

aaagaatacgaatttgctaaagagg

All Databases PubMed Nucleotide Protein Genome Structure OMIM
PMC Journals BooksSearch **PubMed** for **campylobacter acetylglactosaminyltransferase** [Go](#) [Clear](#) [Save Search](#)[About Entrez](#) [Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)[Text Version](#)Display **Summary** Show **20** Sort By [Send to](#)[All: 5](#) [Review: 0](#)

Entrez PubMed

Items 1 - 5 of 5

One page.

[Overview](#)[Help | FAQ](#)[Tutorials](#)[New/Noteworthy](#)[E-Utilities](#)


PubMed Services

[Journals Database](#)[MeSH Database](#)[Single Citation](#)[Matcher](#)[Batch Citation](#)[Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI](#)


Related Resources

[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)


☐ **1** [Blixt O, Vasiliu D, Allin K, Jacobsen N, Warnock D, Razi N, Paulson JC, Bernatchez S, Gilbert M, Wakarchuk W.](#) [Related Articles, Links](#)

 Chemoenzymatic synthesis of 2-azidoethyl-ganglio-oligosaccharides GD3, GT3, GM2, GD2, GT2, GM1, and GD1a. Carbohydr Res. 2005 Sep 5;340(12):1963-72. PMID: 16005859 [PubMed - indexed for MEDLINE]


☐ **2** [Goodfellow JA, Bowes T, Sheikh K, Odaka M, Halstead SK, Humphreys PD, Wagner ER, Yuki N, Furukawa K, Furukawa K, Plomp JJ, Willison HJ.](#) [Related Articles, Links](#)

 Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy. J Neurosci. 2005 Feb 16;25(7):1620-8. PMID: 15716397 [PubMed - indexed for MEDLINE]

☐ **3** [Bowes T, Wagner ER, Boffey J, Nicholl D, Cochrane L, Benboubetra M, Conner J, Furukawa K, Furukawa K, Willison HJ.](#) [Related Articles, Links](#)

 Tolerance to self gangliosides is the major factor restricting the antibody response to lipopolysaccharide core oligosaccharides in Campylobacter jejuni strains associated with Guillain-Barré syndrome. Infect Immun. 2002 Sep;70(9):5008-18. PMID: 12183547 [PubMed - indexed for MEDLINE]

☐ **4** [Guerry P, Szymanski CM, Prendergast MM, Hickey TE, Ewing CP, Pattarini DL, Moran AP.](#) [Related Articles, Links](#)

 Phase variation of Campylobacter jejuni 81-176 lipooligosaccharide affects ganglioside mimicry and invasiveness in vitro. Infect Immun. 2002 Feb;70(2):787-93. PMID: 11796612 [PubMed - indexed for MEDLINE]

☐ **5** [Guerry P, Ewing CP, Hickey TE, Prendergast MM, Moran](#) [Related Articles, Links](#)

AP.



Sialylation of lipooligosaccharide cores affects immunogenicity and serum resistance of *Campylobacter jejuni*.

Infect Immun. 2000 Dec;68(12):6656-62.

PMID: 11083778 [PubMed - indexed for MEDLINE]

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

All Databases PubMed Nucleotide Protein Genome Structure OMIM
PMC Journals BooksSearch for [Save Search](#)[About Entrez](#) [Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)[Text Version](#) Did you mean: *campylobacter acetylgalactosaminyltransferase* (5 items)

Entrez PubMed

[Overview](#)[Help | FAQ](#)[Tutorials](#)[New/Noteworthy](#)[E-Utilities](#)

PubMed Services




[Journals Database](#)[MeSH Database](#)[Single Citation](#)[Matcher](#)[Batch Citation](#)[Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI](#)

Related Resources

[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)Display Show Sort By Send to

Items 1 - 3 of 3

One page.

☐ 1 [Yuki N, Odaka M.](#)[Related Articles, Links](#) [Ganglioside mimicry as a cause of Guillain-Barré syndrome.](#)
Curr Opin Neurol. 2005 Oct;18(5):557-61. Review.
PMID: 16155440 [PubMed - indexed for MEDLINE]☐ 2 [Goodfellow JA, Bowes T, Sheikh K, Odaka M, Halstead
SK, Humphreys PD, Wagner ER, Yuki N, Furukawa K,
Furukawa K, Plomp JJ, Willison HJ.](#)[Related Articles, Links](#) [Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a
antibody-mediated injury in a model of acute motor axonal neuropathy.](#)
J Neurosci. 2005 Feb 16;25(7):1620-8.
PMID: 15716397 [PubMed - indexed for MEDLINE]☐ 3 [Gilbert M, Brisson JR, Karwaski MF, Michniewicz J,
Cunningham AM, Wu Y, Young NM, Wakarchuk WW.](#)[Related Articles, Links](#) [Biosynthesis of ganglioside mimics in Campylobacter jejuni OH4384. Identification of the
glycosyltransferase genes, enzymatic synthesis of model compounds, and characterization
of nanomole amounts by 600-mhz \(1\)h and \(13\)c NMR analysis.](#)
J Biol Chem. 2000 Feb 11;275(6):3896-906.
PMID: 10660542 [PubMed - indexed for MEDLINE]


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

Scholar All articles - **Recent articles** Results 1 - 10 of about 129 for **campylobacter 1,4 acetylgalactosaminyltransferase**

All Results

Did you mean: [campylobacter 14 acetylgalactosaminyltransferase](#)

[M Gilbert](#)

[W Wakarchuk](#)

[N Young](#)

[N Yuki](#)

[M Karwaski](#)

... Variation in the Lipo-oligosaccharide of the Mucosal Pathogen, **Campylobacter jejuni**

[BIOSYNTHESIS OF ... - all 5 versions »](#)

M Gilbert, MF Karwaski, S Bernatchez, NM Young, E ... - Journal of Biological Chemistry, 2002 - ASBMB

... 6, 7). Epidemiological studies have shown that **Campylobacter** infections are ... at 37 °C for 5 min to 2 h. The **-1,4-N-acetylgalactosaminyltransferase** was assayed ...

[Cited by 66](#) - [Related Articles](#) - [Web Search](#)

[Biosynthesis of Ganglioside Mimics in **Campylobacter jejuni** OH4384 IDENTIFICATION OF THE ... - all 3 versions »](#)

M Gilbert, JR Brisson, MF Karwaski, J Michniewicz, ... - Journal of Biological Chemistry, 2000 - ASBMB

... 1). Epidemiological studies have shown that **Campylobacter** infections are more ... for the synthesis of this structure: **-1,4-N-acetylgalactosaminyltransferase**, -1,3 ...

[Cited by 89](#) - [Related Articles](#) - [Web Search](#)

[... of Lipooligosaccharide Cores Affects Immunogenicity and Serum Resistance of **Campylobacter jejuni** - all 6 versions »](#)

P Guerry, CP Ewing, TE Hickey, MM Prendergast, AP ... - Infection and Immunity, 2000 - iai.highwire.org

... of the lipooligosaccharide (LOS) core of **Campylobacter jejuni** MSC57360 ... of a gene encoding a putative **-1,4-N-acetylgalactosaminyltransferase** (Cgt) resulted in ...

[Cited by 44](#) - [Related Articles](#) - [Web Search](#)

[Functional analysis of the **Campylobacter jejuni** N-linked protein glycosylation pathway - all 5 versions »](#)

D Linton, N Dorrell, PG Hitchen, S Amber, AV ... - Molecular Microbiology, 2005 - Blackwell Synergy

... Thus, it has been proposed that the **Campylobacter** N-linked glycan ... infer that the pglJ gene product is an **1,4 N-acetylgalactosaminyltransferase** responsible for ...

[Cited by 23](#) - [Related Articles](#) - [Web Search](#)

[The crucial role of **Campylobacter jejuni** genes in anti-ganglioside antibody induction in Guillain- ... - all 10 versions »](#)

PCR Godschalk, AP Heikema, M Gilbert, T Komagamine ... - Journal of Clinical Investigation, 2004 - Am Soc Clin Investig

... is consistent with the lower activity of the β -**1,4-N-acetylgalactosaminyltransferase** (CgtA) on ... for the biosynthesis of ganglioside mimics in **Campylobacter** LOS ...

[Cited by 43](#) - [Related Articles](#) - [Web Search](#)

[... GlcNAc/Glc 4-Epimerase Supports the Synthesis of Three Cell Surface Glycoconjugates in **Campylobacter** ... - all 5 versions »](#)

S Bernatchez, CM Szymanski, N Ishiyama, J Li, HC ... - Journal of Biological Chemistry, 2005 - ASBMB

Cited by 14 - Related Articles - Web Search

Cited by 73 - Related Articles - Web Search

Cited by 22 - Related Articles - Web Search

Cited by 19 - Related Articles - Web Search

Cited by 27 - Related Articles - Web Search

Result Page: 1 2 3 4 5 6 7 8 9 10 **Next**

campylobacter 1,4 acetyl galactosaminide 4-epimerase

<http://scholar.google.com/scholar?q=campylobacter+1%2C4+acetylgalactosaminyltransferase&hl=en&lr=...> 1/16/2008


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

Scholar All articles - **Recent articles** Results 1 - 10 of about 74 for **campylobacter 1,4 acetylgalactosaminyl trans**

All Results

Did you mean: [campylobacter 14 acetylgalactosaminyltransferase](#)

[M Gilbert](#)
[W Wakarchuk](#)
[R Spiro](#)
[J Brisson](#)
[M Karwaski](#)

Biosynthesis of Ganglioside Mimics in *Campylobacter jejuni* OH4384 IDENTIFICATION OF THE ... - all 3 versions »

M Gilbert, JR Brisson, MF Karwaski, J Michniewicz, ... - Journal of Biological Chemistry, 2000 - ASBMB

... lipooligosaccharide (LOS) of a bacterial pathogen, ***Campylobacter jejuni* OH4384** ... We identified genes encoding a **-1,4-N-acetylgalactosaminyl-transferase (cgtA)**, a ...

[Cited by 89](#) - [Related Articles](#) - [Web Search](#)

Overexpression of GD1a Ganglioside Sensitizes Motor Nerve Terminals to Anti-GD1a Antibody-Mediated ... - all 4 versions »

JA Goodfellow, T Bowes, K Sheikh, M Odaka, SK ... - Journal of Neuroscience, 2005 - neuroscience.org

... by immunizing GD1a-deficient, **-1,4-N-acetylgalactosaminyl transferase** knock-out mice with GD1a ganglioside-mimicking antigens from ***Campylobacter jejuni*** and ...

[Cited by 12](#) - [Related Articles](#) - [Web Search](#)

Carbohydrate mimicry: a new paradigm of autoimmune diseases - all 3 versions »

N Yuki - Current Opinion in Immunology, 2005 - Elsevier

... an oligosaccharide structure (Gal β 1-3 GalNAc β 1-4 [NeuAc α 2-3 ... **Figure 1. Carbohydrate mimicry of GM1/GD1a gangliosides, *Campylobacter jejuni* lipo** ...

[Cited by 15](#) - [Related Articles](#) - [Web Search](#)

Chemoenzymatic synthesis of GM 3 and GM 2 gangliosides containing a truncated ceramide ... - all 2 versions »

S Jacques, JR Rich, CC Ling, DR Bundle - Organic & Biomolecular Chemistry, 2006 - rsc.org

... 2 analogue8(Scheme1).Usingarecombinant b-(1,4)-N**-acetylgalactosaminyl transferase** 6 and UDP-GlcNAc 4- epimerase 17 from ***Campylobacter jejuni***, GM 2 analogue 8 ...

[Cited by 1](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

Nucleic acids encoding β -1, 4-GalNAc transferase - all 3 versions »

M Gilbert, WW Wakarchuk - US Patent 6,911,337, 2005 - Google Patents

... Epidemiological studies have shown that ***Campylobacter*** infections are more ... as shown in SEQ ID NO:1. In presently ... polypeptide that has a (31,4-GalNAc **transferase** ...

[Related Articles](#) - [Web Search](#)

Polypeptides having β -1, 4-GalNAc transferase activity - all 4 versions »

M Gilbert, WW Wakarchuk - US Patent 7,078,207, 2006 - Google Patents

... studies have shown that ***Campylobacter*** infections are ... that has a (3 1 ,4-GalNAc **transferase** ... fructosyl; iucosyl; galactosyl; N**-acetylgalactosaminyl**; glucosyl; N ...

[Related Articles](#) - [Web Search](#)

Ganglioside mimicry as a cause of Guillain-Barré syndrome - all 4 versions »

N Yuki, M Odaka - Curr. Opin. Neurol, 2005 - co-neurology.com

... mimicry of GM1 ganglioside and ***Campylobacter jejuni*** Lipo ... developed standards for

such documentation: (1) It should ... isolates than in enteritis isolates [4,25 ...
[Cited by 9](#) - [Related Articles](#) - [Web Search](#)

Beta 1, 4-N-acetylgalactosaminyltransferases form C. jejuni

M Gilbert, WW Wakarchuk - 2004 - freepatentsonline.com

... Gal=galactosyl; [0050] GalNAc=N-**acetylgalactosaminyl**; [0051] Glc ... Examples of the
.beta.1,4-GalNAc transferases ... that are produced by **Campylobacter** species, such ...

[Cached](#) - [Web Search](#)

[CITATION] Campylobacter jejuni Summary report

S Szczerba, M Pathogenicity

[Related Articles](#) - [Web Search](#)

Complex Gangliosides at the Neuromuscular Junction Are Membrane Receptors for Autoantibodies
and ... - all 4 versions »

RWM Bullens, GM O'Hanlon, E Wagner, PC Molenaar, K ... - Journal of Neuroscience, 2002 -
neuroscience.org

... and originated from MFS/GBS-associated **Campylobacter jejuni** strains ... complement-
inactivated MFS serum (diluted 1:2) or ... procedure comprised a 3-4 hr incubation ...

[Cited by 31](#) - [Related Articles](#) - [Web Search](#)

Did you mean to search for: campylobacter 14 acetylgalactosaminyltransferase

Go o o o o o o o o o g l e ►

Result Page: 1 2 3 4 5 6 7 8 [Next](#)

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google

EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	44	"1,4" adj2(acetylgalactosaminyl transferase)	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:53
L2	60	"1,4" adj2(acetylgalactosaminyltransferase)	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:53
L3	61	l1 or l2	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:53
L4	1	l3 near6 campylobacter	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:54
L5	1	l3 near10 campylobacter	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:54
L6	45	l3 and campylobacter	US-PGPUB; USPAT	ADJ	OFF	2008/01/16 15:54